A GUIDE TO MONITORING SIDE EFFECTS

LEARN WHY IT'S IMPORTANT AND HOW TO KEEP TRACK

WHAT IS IMFINZI?

IMFINZI is a prescription medicine used to treat adults with **a type of lung cancer called SCLC**.

- IMFINZI may be used alone when your LS-SCLC cannot be removed by surgery **and** has responded or stabilized after initial treatment with chemotherapy that contains platinum, given at the same time as radiation therapy.
- IMFINZI may be used with the chemotherapy medicines etoposide and either carboplatin or cisplatin as your first treatment when your SCLC has spread within your lungs or to other parts of the body (extensive-stage small cell lung cancer, or ES-SCLC).

It is not known if IMFINZI is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about IMFINZI® (durvalumab)?

IMFINZI is a medicine that may treat certain cancers by working with your immune system. IMFINZI can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

For Important Safety Information, please see pages 4-8.

Please see additional Important Safety Information throughout and click here for Full Prescribing Information including Medication Guide for IMFINZI.

IMFINZI®
durvalumab
Injection for Intravenous Use 50 mg/mL

Understanding side effects

If you're starting treatment with IMFINZI, it's important for you to understand that you may experience side effects.

IMFINZI is a medicine that may treat certain cancers by working with your immune system.

IMFINZI may cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. There is a possibility that these problems can sometimes become severe or even life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

Monitoring and reporting signs of side effects

Track how you usually feel

You can start keeping track of how you feel before treatment. Taking note of how you normally feel can help you better track any changes that may occur once treatment has started.

Respond to side effects early on

Getting medical treatment right away may help keep any problems you may experience from becoming more serious. Your healthcare provider will check you for side effects during your treatment with IMFINZI. Your healthcare provider may also need to delay or completely stop treatment with IMFINZI if you have severe side effects.

For Important Safety Information, please see pages 4-8.

Stay connected with your care team

It's important to **let your care team know** of any changes in your health. Talk to them about how you're currently feeling, new signs of side effects, or any other change. Remember, a small side effect can lead to something more serious if not treated.

Turn the page for an overview of potential urgent side effects. Use the pages in this brochure to track the signs of side effects throughout your treatment.

If you have any questions about your treatment, ask your doctor. The information in this brochure does not take the place of talking to your care team about your small cell lung cancer (SCLC) treatment.

In case of an emergency, remember to fill out the wallet card in this brochure. Please see page 9.

Side effects to watch out for

Look out for the following side effects so they may be caught and managed early. **Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including:**

Do not attempt to treat any of the side effects without talking to your care team first.

Lung problems

- Cough
- Shortness of breath
- Chest pain

Intestinal problems

- Diarrhea (loose stools) or more frequent bowel movements than usual
- Stools that are black, tarry, sticky, or have blood or mucus
- Severe stomach-area (abdomen) pain or tenderness

Liver problems

- Yellowing of your skin or the whites of your eyes
- Severe nausea or vomiting
- Pain on the right side of your stomach area (abdomen)
- Dark urine (tea colored)
- Bleeding or bruising more easily than normal

 For Important Safety Information, please see pages 6-8.

Hormone gland problems

- Headaches that will not go away or unusual headaches
- Eye sensitivity to light
- Eye problems
- Rapid heartbeat
- Increased sweating
- Extreme tiredness
- Weight gain or weight loss
- Feeling more hungry or thirsty than usual
- Urinating more often than usual
- Hair loss
- Feeling cold
- Constipation
- Your voice gets deeper
- Dizziness or fainting
- Changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

Kidney problems

- Decrease in your amount of urine
- Blood in your urine
- Swelling of your ankles
- Loss of appetite

Skin problems

- Rash
- Itching
- Skin blistering or peeling
- Painful sores or ulcers in mouth or nose, throat, or genital area
- Fever or flu-like symptoms
- Swollen lymph nodes

Pancreas problems

- Pain in your upper stomach area (abdomen)
- Severe nausea or vomiting
- Loss of appetite

Side effects to watch out for (continued)

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with IMFINZI. Call or see your healthcare provider right away for any new or worsening signs or symptoms.

Signs or symptoms may include:

- Chest pain
- Irregular heartbeats
- Shortness of breath or swelling of ankles
- Confusion
- Sleepiness
- Memory problems
- Changes in mood or behavior
- Stiff neck, balance problems; tingling, numbness or weakness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eye sight
- Persistent or severe muscle pain or weakness, muscle cramps, joint pain, joint stiffness or swelling
- Low red blood cells and bruising

Infusion reactions that can sometimes be severe or life-threatening.
Signs and symptoms of infusion reactions may include:

- Chills or shaking
- Itching or rash
- Flushing
- Shortness of breath or wheezing
- Dizziness
- Feel like passing out
- Fever
- Back or neck pain

Rejection of a transplanted organ or tissue. Your healthcare provider should tell you what signs and symptoms you should report and monitor you depending on the type of organ or tissue transplant that you have had

Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with IMFINZI. Your healthcare provider will monitor you for these complications

Additional safety information

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with IMFINZI. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop treatment with IMFINZI if you have severe side effects

Before you receive IMFINZI, tell your healthcare provider about all of your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ or tissue transplant, including corneal transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. IMFINZI can harm your unborn baby

  Females who are able to become pregnant:

  - Your healthcare provider will give you a pregnancy test before you start treatment with IMFINZI.
  - You should use an effective method of birth control during your treatment and for 3 months after the last dose of IMFINZI. Talk to your healthcare provider about birth control methods that you can use during this time.
  - Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with IMFINZI.
- are breastfeeding or plan to breastfeed. It is not known if IMFINZI passes into your breast milk. Do not breastfeed during treatment and for 3 months after the last dose of IMFINZI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of IMFINZI?

IMFINZI can cause serious side effects (see pages 4-6).

The most common side effects of IMFINZI when used alone in adults with limited-stage small cell lung cancer (LS-SCLC) include:

- Inflammation in the lungs
- Feeling tired or weak

The most common side effects of IMFINZI when used with etoposide and either carboplatin or cisplatin medicines in adults with extensive-stage small cell lung cancer (ES-SCLC) include:

- Nausea
- Feeling tired or weak
- Hair loss

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of IMFINZI. Ask your healthcare provider or pharmacist for more information.

Call your doctor for medical advice about side effects. You may report side effects related to AstraZeneca products. If you prefer to report these to the FDA, either visit www.FDA.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and click here for Full Prescribing Information including Medication Guide for IMFINZI.

LS-SCLC side effects tracker

How to track signs of side effects

As you treat limited-stage SCLC, below is a helpful guide you can use to track signs of side effects.

NAME:

	CHANGES IN BREATHING	TIREDNESS
What's normal for me	My breathing disrupts my daily activites A LITTLE A LOT	I usually sleep: ..... hours a day
Date, appointments, etc	Note any other details A LITTLE A LOT	Write down how many hours you've slept last night
1/12 infusion at Dr White	When going up stairs A LITTLE A LOT	I slept 6 hours last night
	A LITTLE A LOT	
	A LITTLE A LOT	
	A LITTLE A LOT	
	A LITTLE A LOT	

NAME:

OTHER SIGNS OF SIDE EFFECTS	WHAT ACTIVITIES DID YOU DO?
I usually experience:	
Write down how often you've noticed this and other signs of side effects today	
	Went to the park today.

LS-SCLC side effects tracker

How to track signs of side effects

As you treat limited-stage SCLC, below is a helpful guide you can use to track signs of side effects.

NAME:

	CHANGES IN BREATHING	TIREDNESS
What's normal for me	My breathing disrupts my daily activites A LITTLE A LOT	I usually sleep: ..... hours a day
Date, appointments, etc	Note any other details A LITTLE A LOT	Write down how many hours you've slept last night
	A LITTLE A LOT	
	A LITTLE A LOT	
	A LITTLE A LOT	
	A LITTLE A LOT	
	A LITTLE A LOT	

NAME:

OTHER SIGNS OF SIDE EFFECTS	WHAT ACTIVITIES DID YOU DO?
I usually experience:	
Write down how often you've noticed this and other signs of side effects today	

LS-SCLC side effects tracker

How to track signs of side effects

As you treat limited-stage SCLC, below is a helpful guide you can use to track signs of side effects.

NAME:

	CHANGES IN BREATHING	TIREDNESS
What's normal for me	My breathing disrupts my daily activites A LITTLE A LOT	I usually sleep: ..... hours a day
Date, appointments, etc	Note any other details A LITTLE A LOT	Write down how many hours you've slept last night
	A LITTLE A LOT	
	A LITTLE A LOT	
	A LITTLE A LOT	
	A LITTLE A LOT	
	A LITTLE A LOT	

NAME:

OTHER SIGNS OF SIDE EFFECTS	WHAT ACTIVITIES DID YOU DO?
I usually experience:	
Write down how often you've noticed this and other signs of side effects today	

ES-SCLC side effects tracker

How to track signs of side effects

As you treat extensive-stage SCLC, below is a helpful guide you can use to track signs of side effects.

NAME:

	NAUSEA	HAIR LOSS	TIREDNESS
What's normal for me	On a scale of 1 to 10, (10 = severe), my nausea is usually:	I've noticed hair shedding or changes to my hair line. ○○○○○ NEVER OFTEN	I usually sleep: ..... hours a day
Date, appointments, etc	Did anything specific trigger your nausea (such as certain foods, smells, activities)?	Write down how often you've noticed hair loss	Write down how many hours you've slept last night
	- 5 - Lasted 3 hours after breakfast	Clumps of hair when brushing ○○X○○ NEVER OFTEN	I slept 6 hours last night
		○○○○○ NEVER OFTEN	
		○○○○○ NEVER OFTEN	
		○○○○○ NEVER OFTEN	
		○○○○○ NEVER OFTEN	

NAME:

OTHER SIGNS OF SIDE EFFECTS	WHAT ACTIVITIES DID YOU DO?
I usually experience:	
Write down how often you've noticed this and other signs of side effects today	
	Went to the park today.

ES-SCLC side effects tracker

How to track signs of side effects

As you treat extensive-stage SCLC, below is a helpful guide you can use to track signs of side effects.

NAME:

	NAUSEA	HAIR LOSS	TIREDNESS
What's normal for me	On a scale of 1 to 10, (10 = severe), my nausea is usually:	I've noticed hair shedding or changes to my hair line. ○○○○○ NEVER OFTEN	I usually sleep: ..... hours a day
Date, appointments, etc	Did anything specific trigger your nausea (such as certain foods, smells, activities)?	Write down how often you've noticed hair loss	Write down how many hours you've slept last night
		○○○○○ NEVER OFTEN	
		○○○○○ NEVER OFTEN	
		○○○○○ NEVER OFTEN	
		○○○○○ NEVER OFTEN	
		○○○○○ NEVER OFTEN	

NAME:

OTHER SIGNS OF SIDE EFFECTS	WHAT ACTIVITIES DID YOU DO?
I usually experience:	
Write down how often you've noticed this and other signs of side effects today	

ES-SCLC side effects tracker

How to track signs of side effects

As you treat extensive-stage SCLC, below is a helpful guide you can use to track signs of side effects.

NAME:

	NAUSEA	HAIR LOSS	TIREDNESS
What's normal for me	On a scale of 1 to 10, (10 = severe), my nausea is usually:	I've noticed hair shedding or changes to my hair line. ○-○-○-○-○ NEVER OFTEN	I usually sleep: ..... hours a day
Date, appointments, etc	Did anything specific trigger your nausea (such as certain foods, smells, activities)?	Write down how often you've noticed hair loss	Write down how many hours you've slept last night
		○-○-○-○-○ NEVER OFTEN	
		○-○-○-○-○ NEVER OFTEN	
		○-○-○-○-○ NEVER OFTEN	
		○-○-○-○-○ NEVER OFTEN	
		○-○-○-○-○ NEVER OFTEN	

NAME:

OTHER SIGNS OF SIDE EFFECTS	WHAT ACTIVITIES DID YOU DO?
I usually experience:	
Write down how often you've noticed this and other signs of side effects today	

Your IMFINZI wallet card

If you ever experience an emergency and need to see a doctor who's not on your regular care team, keeping this card in your wallet can let them know that you're receiving an immunotherapy and help them get in touch with your doctor if they need to.

3 Tips to Make Sure You Get the Appropriate Care

1. Fill out this card and carry it with you at all times.
2. Keep this card with you for at least 3 months, because side effects can still occur during that time.
3. Take a picture of this card with your phone in case you ever leave it at home.

IMFINZI®
durvalumab
Injection for Intravenous Use 50 mg/mL

AstraZeneca
IMFINZI is a registered trademark of the AstraZeneca group of companies.
©2025 AstraZeneca. All rights reserved. US-106872 Last Updated 12/25